



CropAcre PodMix

CropAcre Pulse & Rape Mix is specifically formulated using the unique ACIDplex technology for easy mixing, enhanced compatibility and efficiency of uptake. Using CropAcre PodMix will also improve spraying water quality.

CropAcre PodMix contains:

		% w/w	g/Kg
Manganese	(Mn)	8.00	80
Magnesium	(Mg)	4.00	40.0
	(MgO)	6.4	64.0
Iron	(Fe)	0.25	2.5
Zinc	(Zn)	0.3	3.0
Boron	(B)	4.0	40.0
Molybdenum	(Mo)	0.5	5.0
Sulphur	(S)	10.26	102.6
Equivalent to	(SO ₃)	25.65	256.5

EEC FERTILISER. Use only where there is a recognised need. Do exceed the appropriate application rates.

Rates of Application

Maintenance	1.25 kg/ha
Moderate Deficiency	2.5 kg/ha
Severe Deficiency	2.5 kg/ha Repeated

PRECAUTIONS

USE IN ACCORDANCE WITH COSHH ASSESSMENT WHEN HANDLING OR USING THIS PRODUCT

R-48	HARMFUL. RISK OF SERIOUS DAMAGE TO HEALTH BY PROLONGED EXPOSURE BY INHALATION AND IF SWALLOWED
R51/53	TOXIC TO AQUATIC ORGANISMS. MAY CAUSE LONG TERM DAMAGE IN THE AQUATIC ENVIRONMENT
R-36/38	IRRITATING TO EYES/SKIN
S(2-)	KEEP OUT OF REACH OF CHILDREN

PACK SIZE 10 kg



CROPWELL

Cropwell Ltd
Scalford Station, Melton Mowbray, Leicestershire LE14 4UD
01664444177



CropAcre Products General Recommendations

All products in the CropAcre range are specifically formulated using the uniqueACIDplex technology for easy mixing, enhanced compatibility and efficiency of uptake.

Adding CropAcre products to the spray tank will also improve spraying water quality.

Directions for use:

The full range of field crops, vegetable, fruit and ornamentals may be treated.

Crops responses will be dependent on their susceptibility to any particular deficiency. All edible crops may be treated right up until harvest.

Water Volumes

Do not apply in less than 200 litres of water per hectare unless the spraying equipment is specifically designed for low volume application (e.g. Airtec or low volume hydraulic nozzles).

Timing

As with all trace element sprays frequent repeated applications at relatively low rates are preferable to one large dose. As nutrients are taken in by the foliage sufficient leaf area must be present to absorb the spray. This normally coincides with the 3-leaf stage onwards in most crops. Applications should be made in the early morning or late evening when the uptake by the plant will be maximised.

Mixing

Ensure spray equipment is clean and free of residues from previous use. Half fill the sprayer with water and start agitation. Make sure that the CropAcre product is the first product that is added to the spray tank. This will ensure that the water is suitably conditioned for maximum spraying efficiency. Slowly add the required quantity of product to moving water.

Top up the spray tank with the rest of the water and mix well before spraying.

Do not allow the mixture to stand idle at anytime.

Compatibility

CropAcre products are designed to be physically and chemically compatible with a wide range of agrochemical products. Information is available from your supplier. Variable combinations of factors such as growth stage, seasonal conditions and composition of tank mixes may affect biological compatibility.

DO NOT MIX with hormone weed killers or mixtures containing hormone weed killers unless specific recommendations are given.



CROPWELL

Cropwell Ltd

Scalford Station, Melton Mowbray, Leicestershire LE14 4UD

01664444177